



Short Communication

Present diversity of Galápagos leaf-toed geckos (*Phyllodactylidae*: *Phyllodactylus*) stems from three independent colonization eventsOmar Torres-Carvajal^{a,*}, Andrea Rodríguez-Guerra^a, Jaime A. Chaves^b^a Museo de Zoología, Escuela de Biología, Pontificia Universidad Católica del Ecuador, Avenida 12 de Octubre y Roca, Apartado 17-01-2184, Quito, Ecuador^b Universidad San Francisco de Quito, Colegio de Ciencias Biológicas y Ambientales - Extensión Galápagos, Campus Cumbayá, Casilla Postal 17-1200-841, Quito, Ecuador

ARTICLE INFO

Article history:

Received 11 February 2016

Revised 23 May 2016

Accepted 7 July 2016

Available online 9 July 2016

Keywords:

Galápagos

Island biogeography

Lizards

Multiple colonizations

Phylogeny

Species tree

ABSTRACT

We re-examined the biogeography of the leaf-toed geckos (*Phyllodactylus*) endemic to the Galápagos Islands by analyzing for the first time samples of *P. gilberti*, a species endemic to Wolf island, in a phylogenetic framework. Our aim was to test the three-colonizations scenario previously proposed for these lizards and estimate the age of each colonization event. To achieve this we estimated simultaneously a species tree and divergence times with Bayesian methods. Our results supported the three-colonizations scenario. Similar to a previous hypothesis, the species tree obtained here showed that most species of *Phyllodactylus* are nested in a single clade with an age between 5.49 and 13.8 Ma, whereas a second independent colonization corresponding to *P. darwini* from San Cristóbal island occurred 3.03 Ma ago. The species from Wolf island, *P. gilberti*, stems from a more recent colonization event (0.69 Ma). Thus, present diversity of Galápagos leaf-toed geckos stems from three independent, asynchronous colonization events. As with other Galápagos organisms, the Pacific coast of South America seems to be the source for the founders of *P. gilberti*.

© 2016 Elsevier Inc. All rights reserved.

1. Introduction

Located about 960 km west from the coast of Ecuador, the Galápagos archipelago is widely recognized as one of the best model systems for evolutionary and biogeographic studies (e.g., Emerson, 2002; Grant and Grant, 2014; Losos and Ricklefs, 2009). Such studies have benefited from the relatively well-known geological history of these oceanic islands, which have been forming for millions of years as the Nazca Plate moves eastwards over a volcanic hotspot. As a result, island age increases eastwards along the plate, with the oldest islands—Española, San Cristóbal, Santa Fé—located in the south-eastern part of the archipelago (Geist et al., 2014; White et al., 1993). The northern islands (Genovesa, Marchena, Pinta, Wolf and Darwin), however, are not typical hotspot islands in that they originated from hotspot-derived plume material flowing within the mantle towards the Galápagos Spreading Center, the boundary between the Cocos and Nazca plates (Geist et al., 2014; Harpp and Geist, 2002; Harpp et al., 2002).

Among evolutionary studies of Galápagos land vertebrates, Darwin and Wolf islands have received little attention as they have not been colonized by non-volant land vertebrates, except for Wolf's

leaf-toed geckos, *Phyllodactylus gilberti* Heller 1903. Darwin and Wolf, two of the smallest islands of the archipelago, are located in its remote far north. Darwin has a surface area of 106.3 ha and Wolf is approximately 134.4 ha (Black, 1973). These islands are small caps on mostly submarine volcanoes (Geist et al., 2014). The age of Darwin is about 400 ka, whereas Wolf is between 0.88 ± 0.13 and 1.60 ± 0.07 Ma old (White et al., 1993).

Wolf's leaf-toed geckos represent one of eight named species of endemic leaf-toed geckos known to occur in the Galápagos (*P. barringtonensis* Van Denburgh, 1912; *P. baurii* Garman, 1892; *P. darwini* Taylor, 1942; *P. duncanensis* Van Denburgh 1912; *P. galapagensis* Peters, 1869; *P. gorii* Lanza, 1973; *P. leei* Cope, 1889); in addition, two undescribed species might occur in Isabela and Santiago, respectively (Torres-Carvajal et al., 2014). Based on genetic similarity of allozyme data, Wright (1983) hypothesized that leaf-toed gecko diversity in the Galápagos stems from three colonization events: (i) one that led to the establishment of *P. gilberti* in Wolf; (ii) a second one that led to the establishment of *P. darwini* in San Cristóbal, and (iii) a third one that explains the remaining diversity of leaf-toed geckos in the archipelago, although not necessarily in that order. Wright's hypothesis is very significant for the evolution and biogeography of Galápagos fauna, as other animal radiations in the archipelago stem from a single colonization event, except for lava lizards, which have colonized the islands twice (Benavides et al., 2009; Wright, 1983). In a recent

* Corresponding author.

E-mail address: omartorcar@gmail.com (O. Torres-Carvajal).

study, [Torres-Carvajal et al. \(2014\)](#) confirmed Wright's hypotheses (ii) and (iii) using phylogenetic analyses of DNA sequence data. Here we use genomic samples of *P. gilberti* from Wolf Island to test Wright's hypothesis that this species originated from an independent colonization event. By inferring a dated species tree including all recognized species of Galápagos leaf-toed geckos, we confirm that these reptiles are unique among terrestrial animals in the archipelago in that their origin stems from three independent and asynchronous colonization events.

2. Materials and methods

2.1. Sampling and laboratory protocols

We obtained sequence data of two specimens of *Phyllodactylus gilberti* from Wolf Island (Genbank accession numbers KX268798–KX268815), and combined them with 104 publically available sequences of species of *Phyllodactylus* from Galápagos and continental South America ([Fig. 1](#); [Torres-Carvajal et al., 2014](#)). The gecko *Tarentola mauritanica* (Genbank accession number EU443255) was used as outgroup.

We analyzed 5136 nucleotides (nt) encompassing six nuclear genes and four mitochondrial genes. Nuclear genes included brain-derived neurotrophic factor (BDNF), oocyte maturation factor MOS (CMOS), recombination-activating gene 1 (RAG1), recombination-activating gene 2 (RAG2), acetylcholinergic receptor M4 (ACM4) and phosphatidylinositol 3-kinase (PI3K). Mitochondrial genes included NADH dehydrogenase subunit 4 (ND4), and a continuous fragment including 12S rRNA, tRNA^{Val}, and 16S rRNA.

Total genomic DNA was digested and extracted from muscle tissue (tail tips) using a guanidinium isothiocyanate extraction protocol. Tissue samples were first mixed with Proteinase K and lysis buffer and digested overnight prior to extraction. DNA samples were quantified using a Nanodrop® ND-1000 (NanoDrop Technologies, Inc), re-suspended and diluted to 25 ng/μl in ddH₂O prior to amplification. Primers and amplification protocols are the same as those presented previously ([Torres-Carvajal et al., 2014](#)).

2.2. Phylogenetic analyses

Data were assembled and aligned in Geneious v7 ([Kearse et al., 2012](#)) under default settings for Mafft ([Katoh and Toh, 2010](#)). Ribosomal (12S, tRNA^{Val}, 16S) gene regions with multiple gaps were realigned to minimize indels and optimize nucleotide identities among different individuals. ND4 and nuclear gene sequences were translated into amino acids for confirmation of alignment. We used the same evolutionary models as in [Torres-Carvajal et al. \(2014\)](#).

We estimated a species tree from the multilocus trees under a coalescent model and simultaneously estimated divergence times using ³Beast version 1.8 ([Drummond et al., 2012](#)). Based on previous results ([Torres-Carvajal et al., 2014](#)), we assigned the 107 sequences to 16 species including *P. gilberti*, and calibrated divergence dating using two estimated node ages to set normal priors for the Galápagos main radiation (i.e., the *galapagensis* clade; mean = 5.17 m.a.) and the South American *Phyllodactylus* clade (mean = 23.97 m.a.). The analyses were conducted under a model with uncorrelated substitution rates among branches and the rate for each branch independently drawn from an underlying lognormal distribution ([Drummond et al., 2006](#)); a Yule prior was used for the species tree. To reduce the chance of converging on a local optimum, four runs of 10⁸ generations each were performed. Trees were sampled every 10,000 generations resulting in 10,000 saved trees per analysis. Stationarity was confirmed by plotting the $-\ln L$ per generation in the program Tracer v1.5 ([Rambaut and Drummond, 2007](#)). Independence of samples was assessed by

examining the effective sample sizes (ESS) in Tracer. After analyzing convergence, mixing, and sampling, the first 25% of trees in the sample were discarded as “burn-in” from each run. We then confirmed that the four runs of each Bayesian analysis reached stationarity at a similar likelihood score and that the topologies were similar. We used the resultant trees of each analysis to calculate posterior probabilities (PP) for each bipartition in a maximum clade credibility tree using TreeAnnotator 1.8 ([Rambaut and Drummond, 2016](#)).

3. Results

The species tree ([Fig. 1](#)) was largely concordant with the phylogenetic tree presented by [Torres-Carvajal et al. \(2014; their Fig. 3\)](#); therefore here we present only the results related to *Phyllodactylus gilberti*. This species is sister to *P. reissii* from the Pacific coast in Ecuador and Peru ([Fig. 1](#)) with maximum support (PP = 1). Together they form a clade sister (PP = 0.98) to an unnamed species of *Phyllodactylus* from northern Peru, herein referred to as “*P. reissii*” because of morphological similarity with *P. reissii*. The clade (“*P. reissii*”, (*P. gilberti*, *P. reissii*)) is sister (PP = 1) to a clade containing all species of *Phyllodactylus* (herein named *galapagensis* clade) from Galápagos except *P. darwini*. Thus, among species of *Phyllodactylus* from the Galápagos, *P. gilberti* originated from an independent colonization event.

Similar to the hypothesis presented by [Torres-Carvajal et al. \(2014\)](#), the species tree ([Fig. 1](#)) yielded an estimated node age of 5.49 Ma for the *galapagensis* clade, with a 95% highest posterior density (HPD) interval (i.e., the shortest interval that contains 95% of the sampled values) of 4.54–6.39 Ma, as well as a stem age of 13.8 Ma (HPD = 7.92–20.21 Ma). The split between *P. darwini* from San Cristóbal and its sister Andean species *P. leoni* occurred 3.03 Ma (HPD = 0.22–5.87 Ma), whereas the split between *P. gilberti* and its sister species *P. reissii* was estimated to have occurred 0.69 Ma (HPD = 0.04–1.56 Ma).

4. Discussion

Leaf-toed geckos are unique among terrestrial animals endemic to the Galápagos in that their present diversity is the result of three independent colonizations. Other terrestrial animal taxa in the archipelago stem from a single colonization (see Table 2 in [Torres-Carvajal et al., 2014](#)), except for lava lizards, which have colonized the Galápagos twice ([Benavides et al., 2009](#); [Wright, 1983](#)). A three-colonization scenario was originally proposed for Galápagos leaf-toed geckos by [Wright \(1983\)](#) using genetic distance data, and two of those colonizations were later corroborated by [Torres-Carvajal et al. \(2014\)](#) using DNA-sequence data in a phylogenetic framework. Here we include for the first time samples of Wolf's leaf-toed gecko, *P. gilberti*, and find strong support for [Wright's \(1983\)](#) hypothesis that this species stems from a third colonization event ([Fig. 1](#)). A multiple-colonization scenario for Galápagos geckos is not difficult to explain as these reptiles are capable of travelling long distances by passive drift on ocean currents ([de Queiroz, 2005](#)). Nonetheless, to our knowledge a three-colonization scenario for geckos in other archipelagos of the world has only been proposed for *Tarentola* in the Canary islands ([Carranza et al., 2002](#)).

The Galápagos archipelago is washed by different ocean currents that might represent the invasion routes used by some of the ancestors of its present terrestrial fauna. The Humboldt Current carries cold water northwards along the west coast of South America from the Antarctic to the equator, where it turns westwards to form the South Equatorial Current-South before reaching the Galápagos ([Palacios, 2004](#)). These ocean currents most likely were used




Fig. 1. Top: Map of north-western South America (inset) and the Galápagos Islands showing sampling localities. The 1000-m contour in the Andes is shown in grey on the inset map. Species names of *Phyllodactylus* occurring on each sampled island are provided below each island name. Mainland species include *P. kofordi* (triangles), *P. leoni* (diamond), *P. pumilus* (squares) and *P. reissii* (crosses). Data points for the last species were displaced 2.5 km east for clarity. Bottom: *Phyllodactylus* species tree, with divergence time estimates in millions of years. The species tree is a maximum clade credibility tree of 30,000 samples obtained from a ^{*}Beast analysis of 107 specimens and nine gene regions (5136 nt). The numbers beside branches correspond to posterior probability (PP) values. Node bars represent 95% highest posterior density (HPD) intervals of sampled age-estimate values. Species from the Galápagos Islands (with island of occurrence in parentheses) are in the same colours as the map above. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

as routes for the arrival of the South American ancestors of both *P. darwini*, endemic to San Cristóbal, and the *galapagensis* clade, the only lineage of Galápagos leaf-toed geckos with multiple species, with its most basal species (*P. leei*) also endemic to San Cristóbal (Torres-Carvajal et al., 2014). Given that San Cristóbal is one of the oldest islands and lies on the south-eastern portion of the archipelago, mostly affected by the ocean currents mentioned above, this colonization-route scenario is reasonable for both *P. darwini* and the “*galapagensis*” clade. By contrast, arrival of the ancestor of Wolf’s leaf-toed gecko via the same route implies at least two scenarios that are more complex. First, the ancestor of

P. gilberti could also have arrived to the south-eastern portion of the Galápagos without reaching any of the islands in this area and then dispersed northwest until it reached Wolf island. Alternatively, the ancestor of *P. gilberti* colonized one of the southeastern islands and from there its descendants ‘jumped’ from island to island until they reached and successfully established on Wolf island. The latter scenario implies that ancestors of *P. gilberti* that colonized other islands became extinct, and we have not yet found evidence of those extinctions. Both scenarios raise the question of why other terrestrial animals such as lava lizards and giant tortoises, which also had ancestors arriving to the south-eastern

islands, have not been able to colonize Wolf. Alternatively, arrival of the ancestor of *P. gilberti* to Wolf island could have been influenced by the Panama Current, which carries warm water southwards from the Pacific coast of Central America to the equator and then westwards towards the Galápagos as the South Equatorial Current-North (Houvenaghel, 1984; Palacios, 2004). In fact, Darwin and Wolf Islands are affected mainly by the warm Panama Current (Hearn et al., 2010). Thus, the ancestor of *P. gilberti* could have departed from a population of leaf-toed geckos in the Pacific coast of northwestern South America. Congruent with this idea is the present distribution of *P. reissii*, the sister species of *P. gilberti* (Fig. 1), which occurs along the Pacific coast from northern Peru to northern Ecuador (Dixon and Huey, 1970). In addition, present conditions of both the Panama and South Equatorial Current-North were probably attained 0.69 million years ago, which is our estimate for the split between *P. reissii* and *P. gilberti*. It is less probable that *P. gilberti* has a North or Central American ancestor, as species of *Phyllodactylus* from those areas seem to form a separate clade sister to a South American clade (Torres-Carvajal et al., 2014), where *P. gilberti* is deeply nested. Nonetheless, we recognize that only further sampling of species of *Phyllodactylus* from the continent and other islands (e.g., *P. transversalis* from Malpelo island off the coast from Colombia) will allow a more accurate reconstruction of the colonization routes used by the ancestors of Galápagos leaf-toed geckos.

Even though *P. gilberti* and *P. reissii* diverged recently and are genetically similar (Fig. 1), they have evolved fixed morphological differences. *Phyllodactylus gilberti* differs from *P. reissii* (character states in parentheses) in having a maximum snout-vent length of 55.5 mm (75 mm), small dorsal tubercles (tubercles large, elevated), a median dorsal band of granules distinctly smaller than laterals and usually lighter in color (no medial band of reduced granules, no medial light band), and in lacking tubercles on limbs (tubercles on shanks) (Dixon and Huey, 1970; Van Denburgh, 1912).

Radiations of Galápagos-endemic animals have been estimated to be 1–6 Myr old (Arbogast et al., 2006; Benavides et al., 2009; Caccone et al., 2002; Petren et al., 2005; Schmitz et al., 2007), with the notable exceptions of marine iguanas and *Galapaganus* weevils at one extreme (10–20 Ma and >8.6 Ma, respectively; Rassmann, 1997; Sequeira et al., 2008), and both Galápagos hawks and yellow warblers at the other (<300 kyr; Bollmer et al., 2006; Chaves et al., 2012). Thus, different animal taxa have successfully established in the Galápagos for the last 20 million years following a dynamic and idiosyncratic colonization pattern. The oldest islands in the Galápagos archipelago, Española and San Cristóbal, are thought to have emerged approximately 2–4 Ma (Geist et al., 2014). Therefore, age estimates imply that the ancestors of most Galápagos-endemic vertebrate taxa colonized the archipelago when the present islands were above sea level (Arbogast et al., 2006; Benavides et al., 2009; Caccone et al., 2002; Petren et al., 2005; Wright, 1983). Nonetheless, in agreement with the hypothesis presented by Torres-Carvajal et al. (2014), here we show that the first leaf-toed gecko founders (i.e., ancestors of the *galapagensis* clade) may have landed on what is now submerged island territory sometime between 4.54 and 20.21 Ma. Recent palaeogeographical evidence suggests there were proto-Galápagos islands (now submerged) available for colonization at least 9 Ma (Geist et al., 2014), and although rare, taxa older than the oceanic islands they inhabit also have been documented elsewhere, such as the endemic lobeliads of Hawaii (Givnish et al., 2009). The arrival of the ancestor of *P. darwini* to San Cristóbal 3.03 Ma (HPD = 0.22–5.87 Ma) probably happened when this island already had emerged (i.e., 2.4–4.0 Ma; Geist et al., 2014). In this paper we estimate for the first time the age of Wolf's leaf-toed gecko, *P. gilberti* (Fig. 1; 0.69 Ma; HPD = 0.04–1.56 Ma). Assuming that the ancestor of this species arrived

directly to Wolf island, this probably also happened when this island already had emerged (i.e., 0.88–1.60 Ma; White et al., 1993).

Acknowledgments

We thank D. Anderson, V. Aguirre, F. Ayala, S. Báez, M. Calva, P. Castillo, C. Gaona, F. Gaona, A. León, L. Márquez, S. Ron, I. Tapia, A. Velez, F. Villamar, PUCE students, N. Castillo, and several park rangers of Parque Nacional Galápagos (PNG) for field assistance. Specimens (tail tips) were collected under research permits PC-05-10 and PC-57-15 issued by PNG, Ministerio de Ambiente, for which we thank G. Quezada and W. Tapia. This project was funded by Universidad San Francisco de Quito (GAIAS grant), Pontificia Universidad Católica del Ecuador, Parque Nacional Galápagos and Secretaría de Educación Superior, Ciencia, Tecnología e Innovación.

References

- Arbogast, B.S., Drovetski, S.V., Curry, R.L., Boag, P.T., Seutin, G., Grant, P.R., Grant, B.R., Anderson, D.J., 2006. The origin and diversification of Galapagos mockingbirds. *Evolution* 60, 370–382.
- Benavides, E., Baum, R., Snell, H.M., Sites Jr., J.W., 2009. Island biogeography of Galápagos lava lizards (Tropiduridae: *Microlophus*): species diversity and colonization of the archipelago. *Evolution* 63, 1606–1626.
- Black, J., 1973. Galápagos, archipiélago del Ecuador. Charles Darwin Foundation, Quito, Ecuador.
- Bollmer, J.L., Kimball, R.T., Whiteman, N.K., Sarasola, J.H., Parker, P.G., 2006. Phylogeography of the Galapagos hawk (*Buteo galapagoensis*): a recent arrival to the Galapagos Islands. *Mol. Phylogenet. Evol.* 39, 237–247.
- Caccone, A., Gentile, G., Gibbs, J.P., Fritts, T.H., Snell, H.L., Betts, J., Powell, J.R., 2002. Phylogeography and history of giant Galapagos tortoises. *Evolution* 56, 2052–2066.
- Carranza, S., Arnold, E.N., Mateo, J.A., Geniez, P., 2002. Relationships and evolution of the North African geckos, *Gekkonina* and *Tarentola* (Reptilia: Gekkonidae), based on mitochondrial and nuclear DNA sequences. *Mol. Phylogenet. Evol.* 23, 244–256.
- Chaves, J.A., Parker, P.G., Smith, T.B., 2012. Origin and population history of a recent colonizer, the yellow warbler in Galápagos and Cocos Islands. *J. Evol. Biol.* 25, 509–521.
- de Queiroz, A., 2005. The resurrection of oceanic dispersal in historical biogeography. *Trends Ecol. Evol.* 20, 68–73.
- Dixon, J.R., Huey, R.B., 1970. Systematics of the lizards of the gekkonid genus *Phyllodactylus* of mainland South America. *Los Ang. County Mus. Contrib. Sci.* 192, 1–78.
- Drummond, A.J., Ho, S.Y.W., Phillips, M.J., Rambaut, A., 2006. Relaxed phylogenetics and dating with confidence. *PLoS Biol.* 4, e88.
- Drummond, A.J., Suchard, M.A., Xie, D., Rambaut, A., 2012. Bayesian Phylogenetics with BEAUti and the BEAST 1.7. *Mol. Biol. Evol.* 29, 1969–1973.
- Emerson, B.C., 2002. Evolution on oceanic islands: molecular phylogenetic approaches to understanding pattern and process. *Mol. Ecol.* 11, 951–966.
- Geist, D.J., Snell, H., Snell, H., Goddard, C., Kurz, M.D., 2014. A paleogeographic model of the Galápagos Islands and biogeographical and evolutionary implications. In: Harpp, K.S., Mittelstaedt, E., d'Ozouville, N., Graham, D.W. (Eds.), *The Galápagos: A Natural Laboratory for the Earth Sciences*. John Wiley & Sons, Inc., pp. 145–166.
- Givnish, T.J., Millam, K.C., Mast, A.R., Paterson, T.B., Theim, T.J., Hipp, A.L., Henss, J.M., Smith, J.F., Wood, K.R., Sytsma, K.J., 2009. Origin, adaptive radiation and diversification of the Hawaiian lobeliads (Asterales: Campanulaceae). *Proc. Roy. Soc. B* 276, 407–416.
- Grant, P.R., Grant, B.R., 2014. 40 Years of Evolution: Darwin's Finches on Daphne Major Island. Princeton University Press.
- Harpp, K., Geist, D., 2002. Wolf-Darwin lineament and plume-ridge interaction in northern Galápagos. *Geochim. Geophys. Geosyst.* 3, 1–19.
- Harpp, K.S., Wirth, K.R., Korich, D.J., 2002. Northern Galápagos Province: hotspot-induced, near-ridge volcanism at Genovesa Island. *Geology* 30, 399–402.
- Hearn, A., Ketchum, J., Klimley, A.P., Espinoza, E., Peñaherrera, C., 2010. Hotspots within hotspots? Hammerhead shark movements around Wolf Island, Galapagos Marine Reserve. *Mar. Biol.* 157, 1899–1915.
- Houvenaghel, G.T., 1984. Oceanographic setting of the Galapagos Islands. In: Perry, R. (Ed.), *Key Environments: Galapagos*. Pergamon Press, Oxford, pp. 43–54.
- Katoh, K., Toh, H., 2010. Parallelization of the MAFFT multiple sequence alignment program. *Bioinformatics* 26, 1899–1900.
- Kearse, M., Moir, R., Wilson, A., Stones-Havas, S., Cheung, M., Sturrock, S., Buxton, S., Cooper, A., Markowitz, S., Duran, C., Thierer, T., Ashton, B., Mentjies, P., Drummond, A., 2012. Geneious basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647–1649.
- Losos, J.B., Ricklefs, R.E., 2009. Adaptation and diversification on islands. *Nature* 457, 830–836.

- Palacios, D.M., 2004. Seasonal patterns of sea-surface temperature and ocean color around the Galápagos: regional and local influences. *Deep Sea Res. Part II: Top. Stud. Oceanogr.* 51, 43–57.
- Petren, K., Grant, P.R., Grant, B.R., Keller, L.F., 2005. Comparative landscape genetics and the adaptive radiation of Darwin's finches: the role of peripheral isolation. *Mol. Ecol.* 14, 2943–2957.
- Rambaut, A., Drummond, A.J., 2007. Tracer v1.4. Available at <<http://beast.bio.ed.ac.uk/Tracer>>.
- Rambaut, A., Drummond, A.J., 2016. TreeAnnotator version 1.8.3. Available at <<http://beast.bio.ed.ac.uk>>.
- Rassmann, K., 1997. Evolutionary age of the Galápagos Iguanas predates the age of the present Galápagos Islands. *Mol. Phylogenet. Evol.* 7, 158–172.
- Schmitz, P., Cibois, A., Landry, B., 2007. Molecular phylogeny and dating of an insular endemic moth radiation inferred from mitochondrial and nuclear genes: the genus *Galagete* (Lepidoptera: Autostichidae) of the Galapagos Islands. *Mol. Phylogenet. Evol.* 45, 180–192.
- Sequeira, A.S., Lanteri, A.A., Albelo, L.R., Bhattacharya, S., Sijapati, M., 2008. Colonization history, ecological shifts and diversification in the evolution of endemic Galapagos weevils. *Mol. Ecol.* 17, 1089–1107.
- Torres-Carvajal, O., Barnes, C.W., Pozo-Andrade, M.J., Tapia, W., Nicholls, G., 2014. Older than the islands: origin and diversification of Galápagos leaf-toed geckos (Phyllodactylidae: *Phyllodactylus*) by multiple colonizations. *J. Biogeogr.* 41, 1883–1894.
- Van Denburgh, J., 1912. Expedition of the California Academy of Sciences to the Galapagos Islands, 1905–1906. VI. The geckos of the Galapagos Archipelago. *Proc. Calif. Acad. Sci.* 1, 405–430.
- White, W.M., McBirney, A.R., Duncan, R.A., 1993. Petrology and geochemistry of the Galápagos Islands: portrait of a pathological mantle. *J. Geophys. Res.* 98, 533–563.
- Wright, J.W., 1983. The evolution and biogeography of the lizards of the Galápagos Archipelago: evolutionary genetics of *Phyllodactylus* and *Tropidurus* populations. In: Bowman, R.I., Berson, M., Levinton, A.E. (Eds.), *Patterns of Evolution in Galápagos Organisms*. American Association for the Advancement of Science, Pacific Division, San Francisco, CA, pp. 123–155.